



EUROPLANT

JULINKA



early, mainly firm cooking ware potato

- ✓ yellow flesh colour
- ✓ early high marketable yield
- ✓ smooth skin, shallow eyes
- ✓ early good washability

JULINKA - washed, brushed or fresh, always a good solution

Growing recommendations				
Site requirements		prefers average to better soil conditions with uniform supply of water and nutrients		
Planting 	Seed material sorting 35/55	Row width		Tubers/ha
		75 cm	90 cm	
		28-30 cm	23-25 cm	46,000
	Planting depth	planted 2 cm deeper for high-yielding locations (max. 17 cm soil cover)		
	Seed preparation	has a good dormancy; sprout stimulation evenly before planting (7-10 days)		
	Seed dressing	dressing against Rhizoctonia solani recommended		
Herbicides		no sensitivities are known when used properly		
Foliage		average leaf health; timely and regular use of fungicides against late blight is required		
Harvest		ensure gentle harvest		
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)		medium - compared to the local standard (2/3 before planting and 1/3 later)		
Phosphorus (P ₂ O ₅)		100 kg/ha (water soluble phosphate at planting)		
Potassium (K ₂ O)		350-270 kg/ha (up to 150 kg/ha chloride possible; on lighter soils split doses)		
Magnesium (MgO)		70 kg/ha (for yield and quality assurance)		
Variety characteristics				
Maturity	early	Tuber characteristics Tuber shape Eye depth Flesh colour Skin colour Skin Number of tubers	oval shallow yellow yellow smooth medium	
Cooking type	B			
Consumption quality	mainly firm cooking; very stable in colour			
Yield	medium to high yield in a tight packing grading	Resistances PCN	Ro1 (9), Ro4 (9)	
Youth development	initially hesitant; later rapid	Sensitivities Common scab Spraing Bruising Mechanical damage	low low low low - medium	
Dormancy	good dormancy			

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)